

# SUPPLEMENTAL LABELING FOR USE ON MINT TOUGH® 5 EC HERBICIDE

EPA Reg. No. 91746-5

This Supplemental Label expires on April 14, 2029, and must not be used or distributed after this date.


Active Ingredient:

Pyridate* (Carbonothioic acid, O-(6-chloro-3-phenyl-4-pyridiziny) S-octyl ester): .....	56.07%
Other Ingredients: .....	43.93%
Total:	100.0%

\*Contains 5 lbs. of pyridate per gallon of product.

**KEEP OUT OF REACH OF CHILDREN**

## WARNING



It is a violation of federal law to use this product in a manner inconsistent with its labeling. The labeling must be in possession of the user at the time of application.

Read the label affixed to the container for Tough® 5 EC Herbicide before applying.

Use of Tough® 5 EC according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Tough® 5 EC.

## DIRECTIONS FOR USE

---

### MINT

---

Tough® 5 EC can be applied by ground application and as a broadcast postemergence treatment to mint with a pre-harvest interval of 30 days.

Apply Tough® 5 EC at 24 fl. oz./A per application. Two applications of Tough® 5 EC at 24 fl.oz./A can be made but do not exceed a total of 48 fl. oz. per treated acre per year. Do not exceed a total of 24 fl. oz./A per application.

For a list of weeds controlled by an application of Tough® 5 EC see Table 1. Application of Tough® 5 EC at rates less than 24 fl. oz./A (0.94 lb a.i./A) in postemergence may result in incomplete weed control.

For best results, apply Tough® 5 EC to actively growing weeds and when weeds are small (before the 4-leaf stage). Weeds which emerge after application will not be controlled by Tough® 5 EC.

Always add an appropriate adjuvant to the spray tank (see the SPRAY ADDITIVES section of this label).

Optimal weed control is obtained when Tough® 5 EC is used in a weed control program following a broad spectrum preemergence herbicide. Refer to individual product labels for precautionary statements, restrictions, rates, approved uses and a list of weeds controlled. Add sethoxydim (e.g. Poast® Herbicide, EPA Reg. No. 7969-58), clethodim (e.g. Select Max® Herbicide, EPA Reg. No. 59639-132) or quizalofop-P-ethyl (e.g. Dupont™ Assure® II Herbicide, EPA Reg. No. 352-541) for postemergence grass weed control. Tough® 5 EC may be tank mixed with other registered herbicides for control of additional broadleaf weeds and/or grasses.

Refer to individual product labels for precautionary statements, restrictions, rates, approved uses and a list of weeds controlled.

### **RESTRICTIONS**

- Do not apply more than a total of 48 fl. oz (1.88 lb. pyridate active ingredient) of Tough® 5 EC per acre per year.
- Do not exceed 24 fl. oz. (0.94 lb. a.i./A) in a single postemergence application.
- To avoid excess residues, do not apply within 30 days of harvest
- Re-treatment interval (RTI): Allow at least 14 days between applications of this product.
- DO NOT apply this product through any type of irrigation equipment
- DO NOT apply this product by aerial application
- DO NOT apply this product when wind conditions will allow drift to adjacent susceptible vegetation

Produced for:  
Certis Belchim B.V.  
Stadsplateau 16, Box 607  
3500 AP Utrecht  
Netherlands  
EPA Reg. No. 91746-5  
EPA Establishment No.: \_\_\_\_\_  
Made in \_\_\_\_\_



Distributed By:  
Belchim Crop Protection US Corporation  
225 Wilmington West  
Chester Pike, Suite 200  
Chadds Ford, PA 19317  
Phone: 855-445-7990  
Info.usa@belchim.com